

Cite as: C. M. Minutti et al., *Science*
10.1126/science.aaj20067 (2017).

Local amplifiers of IL-4R α -mediated macrophage activation promote repair in lung and liver

Carlos M. Minutti,^{1,2,3} Lucy H. Jackson-Jones,^{3*} Belén García-Fojeda,^{1,2*} Johanna A. Knipper,³ Tara E. Sutherland,^{3,4} Nicola Logan,³ Emma Rinqvist,^{3†} Raquel Guillamat-Prats,^{2,5} David A. Ferencik,³ Antonio Artigas,^{2,5} Cordula Stamme,⁶ Zissis C. Chronos,⁷ Dietmar M. Zaiss,³ Cristina Casals,^{1,2‡} Judith E. Allen^{3,8‡}

¹Department of Biochemistry and Molecular Biology I, Complutense University of Madrid, 28040-Madrid, Spain. ²Centro de Investigación Biomédica en Red de Enfermedades Respiratorias (CIBERES), Instituto de Salud Carlos III, 28029-Madrid, Spain. ³School of Biological Sciences and School of Clinical Sciences, University of Edinburgh, Edinburgh EH9 3FL, UK. ⁴Faculty of Biology, Medicine and Health, Manchester Collaborative Centre for Inflammation Research, University of Manchester, Manchester M13 9NT, UK. ⁵Critical Care Centre, Corporació Sanitària Universitària Parc Taulí, Universitat Autònoma de Barcelona Parc Taulí 1, 08208-Sabadell, Spain. ⁶Division of Cellular Pneumology, Research Center Borstel, Leibniz Center for Medicine and Biosciences 23845, Borstel and Department of Anesthesiology and Intensive Care, University of Lübeck, 23538 Lübeck, Germany. ⁷Pulmonary Immunology and Physiology Laboratory, Department of Pediatrics, and Microbiology and Immunology, The Pennsylvania State University, College of Medicine, Hershey PA 17033, USA. ⁸Faculty of Biology, Medicine and Health, Wellcome Centre for Cell-Matrix Research, Manchester Academic Health Science Centre, University of Manchester, Manchester M13 9PT, UK.

*These authors contributed equally to this work. †Present address: Department of Medicine Solna, Karolinska Institutet, 17176 Stockholm, Sweden.

‡Corresponding authors. Email: judi.allen@manchester.ac.uk (J.E.A.); ccasals@ucm.es (C.C.)

The type 2 immune response controls helminth infection and maintains tissue homeostasis but can lead to allergy and fibrosis if not adequately regulated. We have discovered local tissue-specific amplifiers of type-2 mediated-macrophage activation. In the lung, surfactant protein A (SP-A) enhanced IL-4-dependent macrophage proliferation and activation, accelerating parasite clearance and reducing pulmonary injury following infection with a lung-migrating helminth. In the peritoneal cavity and liver, C1q enhancement of type 2 macrophage activation was required for liver repair following bacterial infection, but resulted in fibrosis following peritoneal dialysis. IL-4 drives production of these structurally related defense collagens, SP-A and C1q, and the expression of their receptor, myosin 18A. These findings reveal the existence within different tissues of an amplification system needed for local type 2 responses.

The type 2 cytokines IL-4 and IL-13, which signal through IL-4R α , trigger a specialized macrophage phenotype (M(IL-4)) (1) that promotes control of helminth infection (2) and tissue repair (3, 4). M(IL-4)s also contribute to pathology associated with type 2 immunity, including allergy, asthma, and fibrosis (4). However, little is known about tissue-specific factors that might promote both beneficial and detrimental actions of M(IL-4)s.

In the lung, alveolar macrophages (aM ϕ s), together with the respiratory epithelium, are covered by pulmonary surfactant, a lipid-protein network in which surfactant protein A (SP-A) constitutes the major protein component (5). SP-A is a versatile recognition protein (5) that is a member of a group of secreted soluble defense collagens that include the first component of the complement system (C1q), collectins (e.g., SP-A, SP-D, mannan-binding lectin), ficolins, and adiponectin (6). Because of its abundance and known role in immune defense (5), we asked whether SP-A was involved in the local regulation of M(IL-4) effector function in the lung.

M(IL-4)s have a critical role in lung repair following infection with the lung-migrating nematode *Nippostrongylus brasiliensis* (3). We therefore infected WT, IL-4R α -deficient and SP-A-deficient mice with *N. brasiliensis* infective larvae.

Larvae migrate to the lung where they mature for ~2 days, and reach the small intestine by 3 days post infection (p.i.). The type 2 response peaks 6-7 days after inoculation. We observed an up-regulation of SP-A protein (Fig. 1A) and mRNA (fig. S1A) in lungs of *N. brasiliensis* infected C57BL/6 mice at 6 days p.i., which was dependent on IL-4R α . Consistent with a role for SP-A during type 2 immunity to nematode infection, SP-A-deficient mice had greater adult worm burdens (Fig. 1B) and egg output (fig. S1B), and significantly impaired lung repair processes (Fig. 1, C and D) than wild type (WT) mice. The failure to heal was associated with a failure to up-regulate tissue-repair related gene *Colla1* (Collagen, type I, alpha 1) (Fig. 1E) and increased expression of *Mmp12*, an extracellular matrix-degrading enzyme (Fig. 1F). Greater lung damage in SP-A-deficient mice was indicated by increased numbers of red blood cells and neutrophils in bronchoalveolar lavage (BAL) at 6 days p.i (Fig. 1, G and H). The absence of SP-A resulted in reduced expression of the M(IL-4) proteins RELM α (Fig. 1I), Ym1, and arginase (fig. S1, C and D) in aM ϕ s. Secretion of RELM α and Ym1 protein into the alveolar fluid (fig. S1E) was also reduced in SP-A-deficient mice compared with WT mice.

Consistent with the known ability of IL-4 to cause mac-

rophage proliferation during helminth infection (7), aM ϕ s from WT mice exhibited significant proliferation (Fig. 1, J and K) and increased aM ϕ s numbers (fig. S1F) 6 days following *N. brasiliensis* infection. However, SP-A-deficient mice failed to exhibit significantly enhanced aM ϕ proliferation (Fig. 1, J and K) resulting in fewer total macrophage numbers relative to WT mice (fig. S1F). There was no evidence that differences were caused by a failure of SP-A-deficient mice to mount appropriate ILC2, T_H2 (fig. S1, G and H) or local type 2 cytokine responses (fig. S1I). Importantly, aM ϕ s from uninfected SP-A-deficient mice were normal in number, phenotype and ability to respond to IL-4 ex vivo (fig. S2, A to D).

To ascertain if defects in SP-A-deficient mice were caused by defective IL-4R α responsiveness in vivo, WT and SP-A-deficient mice were injected intra-peritoneally (ip) with IL-4 complex (IL-4c). IL-4c delivery increased the amount of SP-A protein in BAL (Fig. 1L) in WT mice and induced M(IL-4) markers and proliferation in aM ϕ s isolated from BAL (Fig. 1, M and N, and fig. S3, A to C) of WT but not SP-A-deficient mice, reflected by diminished secretion of RELM α and Ym1 to the alveolar fluid in SP-A-deficient mice (fig. S3D). IL-4c delivery decreased IL-4R α expression in both WT and SP-A-deficient aM ϕ s (fig. S3E).

Enhancement of M(IL-4)s may underlie the previously reported contribution of SP-A to tissue integrity in other models of acute lung injury (8, 9). The pro-type 2 effects we report here contrast with reports that associate SP-A with protection in asthma (10). However, in addition to promoting M(IL-4) and proliferation of macrophages, the anti-inflammatory properties of SP-A (5, 6, 8, 9, 11) may suppress the strong inflammatory responses that are responsible for more severe asthma. Our data are supported by the finding that SP-D-deficient mice, which lack SP-A (12), also have reduced M(IL-4) responsiveness (13).

To determine whether SP-A acts directly on aM ϕ s, we first tested the ability of adherence-purified macrophages from the alveolar and peritoneal spaces to proliferate in vitro in response to 1 μ g/ml of IL-4; aM ϕ exhibited significant proliferation, but peritoneal macrophages (pM ϕ) failed to proliferate despite expressing M(IL-4) activation markers (fig. S4A). We then tested the ability of SP-A to enhance IL-4 treatment and included C1q as a control because it is a defense collagen structurally homologous to SP-A (5, 6). We found that SP-A, but not C1q, significantly boosted IL-4-mediated aM ϕ proliferation and M(IL-4) markers (Fig. 2A). IL-4R α -deficient aM ϕ showed no proliferation or activation when stimulated with IL-4 (1 μ g/ml) in the absence or presence of SP-A (fig. S4B). Notably, SP-A significantly enhanced proliferation and activation induced by IL-4 in both human (fig. S4C) and rat aM ϕ s (fig. S4D).

To our surprise C1q, but not SP-A, significantly increased

IL-4-mediated proliferation and M(IL-4) marker expression in pM ϕ (Fig. 2A). To verify these findings in vivo, IL-4c was delivered to C1qa-deficient mice. Consistent with the in vitro studies, mice lacking C1q exhibited reduced IL-4-dependent activation and proliferation in pM ϕ but not aM ϕ (Fig. 2B). Similar to SP-A in the lung (Fig. 1L), C1q levels increased in the peritoneal fluid after IL-4c delivery (fig. S5A), indicating that IL-4 drives production of a local signal to amplify its effect on tissue macrophages. Importantly, the pM ϕ phenotype, number, and ability to respond to IL-4 ex vivo are normal in C1qa-deficient mice (fig. S5, B to F). Thus, SP-A and C1q were induced by IL-4 in the lung and peritoneal cavity, respectively, where they acted to enhance proliferation and M(IL-4) activation in a tissue-specific manner.

To determine which receptor mediates SP-A effects on IL-4-stimulated aM ϕ s, we inhibited known receptors for SP-A (5). We observed that the blockade of Myosin18A (Myo18A, *aka* SP-R210), but not signal inhibitory regulatory protein α (SIRP α) or calreticulin (*aka* cC1qR), abrogated SP-A-mediated enhancement of IL-4-induced arginase activity in rat aM ϕ s (fig. S6A). Blockade or RNA silencing of Myo18A consistently abolished SP-A-mediated enhancement of IL-4-induced proliferation and activation of mouse (Fig. 2C), human (fig. S6B) and rat (fig. S6C) aM ϕ s. Myo18A is an unconventional myosin that does not operate as a traditional molecular motor, having both intracellular and cell-surface locations (14), and was recently defined as CD245 (15). Immune activation results in Myo18A localization on the cell surface, where it binds to the collagen-like domain of SP-A (6, 16) and we confirmed that an intact collagen-like domain is required to enhance IL-4-mediated type 2 responses (fig. S6D).

Because C1q is structurally homologous to SP-A in its supra-trimeric assembly and collagen tail (5, 6), we addressed whether Myo18A was also responsible for the actions of C1q. Indeed, blockade of Myo18A prevented C1q enhancement of IL-4 driven activation and proliferation of pM ϕ (Fig. 2C). In vitro, IL-4 promoted Myo18A localization on the cell surface of both aM ϕ and pM ϕ (fig. S7, A and B), which was maximal 24 hours after stimulation. Cell surface expression of Myo18A was similarly observed in vivo following IL-4c delivery (fig. S7, C and D), independent of the presence or absence of SP-A or C1q. We confirmed the role of Myo18A in vivo by intranasal or intraperitoneal delivery of anti-Myo18A antibody. Receptor blockade significantly reduced IL-4-induced proliferation and M(IL-4) activation of aM ϕ and pM ϕ (Fig. 2D), as well as secretion of RELM α and Ym1 to the alveolar and peritoneal fluid (fig. S7, E and F). Thus, Myo18A receptor blockade in the lung or peritoneal cavity phenocopied SP-A or C1qa deficiency, respectively. Together these data indicate that Myo18A is a common receptor or co-receptor for defense collagens present on aM ϕ s and

pM ϕ s, which determines macrophage capacity to respond to IL-4 and whose cell surface expression is itself induced by IL-4. Myo18A lacks a trans-membrane domain (14), and thus must act in concert with transmembrane co-receptors for signal transduction that likely impart tissue-specificity.

The relevance of C1q as a local factor to enhance type 2 responses is supported by the fact that unlike most other complement components, the majority of C1q is produced by myeloid cells in peripheral tissues (17). To determine the physiological relevance of C1q in this context, we evaluated its role in a murine model of peritoneal fibrosis (18). Peritoneal fibrosis is a frequent and serious consequence of peritoneal dialysis (19) associated with alternatively activated macrophages in both humans and mice (18–20). We administered Dianeal-PD4, a clinically-used lactate-based dialysate, every other day for 28 days to WT, C1qa-, and IL-4R α -deficient mice. In WT, but not in C1qa-deficient mice, Dianeal-PD4 treatment provoked the induction of C1q (Fig. 3A) and morphologic changes in tissue sections of the parietal peritoneum, showing significant enlargement of the submesothelial zone caused by collagen deposition (Fig. 3, B and C). Dianeal-PD4 treatment induced markers of fibrosis including collagen mRNAs (*Col1a1* and *Col3a1*) (Fig. 3D and fig. S8A), alpha-smooth muscle actin (*Acta2*) (Fig. 3E), and vascular endothelial growth factor (*Vegf*) (fig. S8B). Significant upregulation of these markers was not observed in C1qa-deficient mice. Conversely, *Mmp12* mRNA was upregulated in C1qa-deficient mice relative to WT mice (fig. S8C) consistent with monocyte infiltration (fig. S8D) (21), and an anti-inflammatory role for C1q (22). Dianeal-PD4 treatment also induced intracellular expression and protein secretion of the M(IL-4) markers RELM α , Ym1, and Arg (Fig. 3, F to H) and moderate proliferation of pM ϕ s (Fig. 3I) in WT but not C1qa-deficient mice. Despite the clear induction of M(IL-4) markers by PD4 delivery and their dependence on C1q, responses of IL-4R α -deficient mice were equivalent to WT mice (Fig. 3, A to I, and fig. S8, A to D). Although initially surprising, Dianeal-PD4 is a lactate-based solution, and lactate can induce M(IL-4) markers by acting downstream of IL-4R α through stabilization of HIF1 α protein (23). Indeed, we observed that the induction of proliferation and M(IL-4) markers following Dianeal-PD4 delivery was essentially absent in mice lacking HIF1 α in macrophages (fig. S9). Together, our data indicate that C1q significantly amplifies peritoneal fibrosis by promoting a type 2 macrophage phenotype driven by lactate and dependent on HIF1 α . These results are consistent with human studies in which C1q is strongly associated with increased fibrosis of skeletal muscle (24).

Critically, we needed to ascertain whether C1q functioned as a type 2 amplifier in tissues beyond the peritoneal cavity and settings that are dependent on IL-4R α . We thus

assessed Myo18A expression on the cell surface of resident macrophages from mice treated with or without IL-4c (Fig. 4A). Consistent with our functional data, Myo18A was expressed in macrophages from the lung and peritoneal cavity, as well as in liver, spleen and adipose tissue, and significantly increased by IL-4 exposure. In contrast, there was minimal Myo18A on pleural cavity macrophages, which explained our failure to identify a role for C1q in the pleural cavity following IL-4c delivery (fig. S10). As predicted by the Myo18 expression data, we found that macrophages from the liver of C1qa-deficient mice had significantly lower levels of proliferation and M(IL-4) activation compared with WT mice following IL-4c delivery (Fig. 4B). Importantly, we observed C1q upregulation in the liver of IL-4c treated mice (Fig. 4C). Of relevance, the number and phenotype of liver macrophages are normal in C1qa-deficient mice (fig. S11). Antibody blockade of Myo18A following IL-4 treatment of isolated liver macrophages verified that C1q generated its effects through Myo18A (fig. S12). We thus sought a model in which M(IL-4)s in the liver played a substantive role.

Infection of the liver by the Gram-positive bacteria *Listeria monocytogenes* causes necroptotic death of resident liver macrophages (Kupffer cells) followed by recruitment of monocytes, which control *L. monocytogenes* infection (25) and repopulate the liver macrophage population (26). Following an initial type 1 response, the type 2 response begins at 3 days p.i. with IL-4-mediated activation and proliferation of liver macrophages acting to repair infection damage (26). To ascertain the relevance of C1q and confirm the role of IL-4R α in this process, we infected WT, C1qa- and IL-4R α -deficient mice with *L. monocytogenes* and performed sample analysis 3.5 days later. Infection resulted in IL-4R α -dependent up-regulation of C1q mRNA in liver (Fig. 4C) while enhanced expression of RELM α and Ym1 by liver macrophages (Fig. 4D and fig. S13A) and macrophage proliferation (Fig. 4E and fig. S13B) were dependent on both IL-4R α and C1q. Consistent with the requirement for basophil-derived IL-4 in *L. monocytogenes*-induced liver macrophage proliferation (26), we observed increases in IL-4 and IL-13 cytokines in liver homogenates (fig. S13C). C1q and IL-4R α deficiency were associated with increased liver injury as assessed by liver transaminases in blood (Fig. 4F) and a failure to up-regulate tissue-repair related genes (*Acta2* and *Col1a1*) (Fig. 4G). Notably, at 3.5 days p.i., liver bacterial burden was higher in WT than in C1qa- or IL-4R α -deficient mice (Fig. 4H). Gene-deficient mice had increased numbers of recruited monocytes (Fig. 4I) and higher iNOS expression in monocytes (CD11b⁺ Ly6C⁺) and liver macrophages (CD11b⁺ F4/80⁺) (fig. S13, D and E) but pro-inflammatory cytokines in liver homogenates were comparable (fig. S13F). These data suggest that C1q, through its ability to orchestrate IL-4R α -dependent type-2-mediated responses, de-

increases the bactericidal capacity of monocyte-derived macrophages, but also limits liver injury and promotes the return to homeostasis (Fig. 4, D to I, and fig. S13).

M(IL-4)s have recently emerged as important players in homeostatic processes (27), but IL-4R α -dependent pathways are amplified during helminth infection with uncontrolled amplification leading to fibrosis (4, 28, 29). Our data show that IL-4 drives production of local specific factors (SP-A and C1q) and expression of their receptor (Myo18A) on the macrophage surface for full M(IL-4) activation and proliferation (fig. S14). These findings reveal the existence within distinct tissues of an amplification system needed for type 2 function. SP-A and C1q are typically produced by alveolar epithelial type II and myeloid cells, respectively, indicating that several different cell types must respond to IL-4 for signal amplification.

The study raises a number of critical questions. What are the Myo18A co-receptors that mediate tissue-specificity? What factors negatively regulate or stop the positive M(IL-4) loop? What are the intracellular signaling pathways regulated by Myo18A and its co-receptors? Critically, SP-A, C1q, and Myo18A are highly conserved across mammalian species, and we have shown the ability of SP-A and Myo18A to enhance human alveolar M(IL-4) proliferation.

REFERENCES AND NOTES

1. P. J. Murray, J. E. Allen, S. K. Biswas, E. A. Fisher, D. W. Gilroy, S. Goerdt, S. Gordon, J. A. Hamilton, L. B. Ivashkiv, T. Lawrence, M. Locati, A. Mantovani, F. O. Martinez, J.-L. Mege, D. M. Mosser, G. Natoli, J. P. Saeij, J. L. Schultze, K. A. Shirey, A. Sica, J. Suttles, I. Udalova, J. A. van Ginderachter, S. N. Vogel, T. A. Wynn, Macrophage activation and polarization: Nomenclature and experimental guidelines. *Immunity* **41**, 14–20 (2014). doi:10.1016/j.immuni.2014.06.008 Medline
2. R. K. Grencis, Immunity to helminths: Resistance, regulation, and susceptibility to gastrointestinal nematodes. *Annu. Rev. Immunol.* **33**, 201–225 (2015). doi:10.1146/annurev-immunol-032713-120218 Medline
3. F. Chen, Z. Liu, W. Wu, C. Rozo, S. Bowdridge, A. Millman, N. Van Rooijen, J. F. Urban Jr., T. A. Wynn, W. C. Gause, An essential role for TH2-type responses in limiting acute tissue damage during experimental helminth infection. *Nat. Med.* **18**, 260–266 (2012). doi:10.1038/nm.2628 Medline
4. T. A. Wynn, K. M. Vannella, Macrophages in Tissue Repair, Regeneration, and Fibrosis. *Immunity* **44**, 450–462 (2016). doi:10.1016/j.immuni.2016.02.015 Medline
5. J. R. Wright, Immunoregulatory functions of surfactant proteins. *Nat. Rev. Immunol.* **5**, 58–68 (2005). doi:10.1038/nri1528 Medline
6. A. J. Tenner, Membrane receptors for soluble defense collagens. *Curr. Opin. Immunol.* **11**, 34–41 (1999). doi:10.1016/S0952-7915(99)80007-7 Medline
7. D. Rückerl, J. E. Allen, Macrophage proliferation, provenance, and plasticity in macroparasite infection. *Immunol. Rev.* **262**, 113–133 (2014). doi:10.1111/imr.12221 Medline
8. H. Goto, J. G. Ledford, S. Mukherjee, P. W. Noble, K. L. Williams, J. R. Wright, The role of surfactant protein A in bleomycin-induced acute lung injury. *Am. J. Respir. Crit. Care Med.* **181**, 1336–1344 (2010). doi:10.1164/rccm.200907-10020C Medline
9. E. N. Atochina, J. M. Beck, A. M. Preston, A. Haczku, Y. Tomer, S. T. Scanlon, T. Fusaro, J. Casey, S. Hawgood, A. J. Gow, M. F. Beers, Enhanced lung injury and delayed clearance of *Pneumocystis carinii* in surfactant protein A-deficient mice: Attenuation of cytokine responses and reactive oxygen-nitrogen species. *Infect. Immun.* **72**, 6002–6011 (2004). doi:10.1128/IAI.72.10.6002-6011.2004 Medline
10. J. G. Ledford, A. M. Pastva, J. R. Wright, Review: Collectins link innate and adaptive immunity in allergic airway disease. *Innate Immun.* **16**, 183–190 (2010). doi:10.1177/1753425910368446 Medline
11. C. M. Minutti, B. García-Fojeda, A. Sáenz, M. de Las Casas-Engel, R. Guillamat-Prats, A. de Lorenzo, A. Serrano-Mollar, Á. L. Corbí, C. Casals, Surfactant Protein A Prevents IFN- γ /IFN- γ Receptor Interaction and Attenuates Classical Activation of Human Alveolar Macrophages. *J. Immunol.* **197**, 590–598 (2016). doi:10.4049/jimmunol.1501032 Medline
12. M. Ikegami, J. A. Whitsett, A. Jobe, G. Ross, J. Fisher, T. Korfhagen, Surfactant metabolism in SP-D gene-targeted mice. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **279**, L468–L476 (2000). Medline
13. S. Thawer, J. Auret, C. Schnoeller, A. Chetty, K. Smith, M. Darby, L. Roberts, R.-M. Mackay, H. J. Whitwell, J. F. Timms, J. Madsen, M. E. Selkirk, F. Brombacher, H. W. Clark, W. G. C. Horsnell, Surfactant Protein-D Is Essential for Immunity to Helminth Infection. *PLOS Pathog.* **12**, e1005461 (2016). doi:10.1371/journal.ppat.1005461 Medline
14. S. Guzik-Lendrum, S. M. Heissler, N. Billington, Y. Takagi, Y. Yang, P. J. Knight, E. Homsher, J. R. Sellers, Mammalian myosin-18A, a highly divergent myosin. *J. Biol. Chem.* **288**, 9532–9548 (2013). doi:10.1074/jbc.M112.441238 Medline
15. A. De Masson, J. Giustiniani, A. Marie-Cardine, J. D. Bouaziz, N. Dulphy, D. Gossot, P. Validire, A. Tazi, C. Garbar, M. Bagot, Y. Merrouche, A. Bensussan, Identification of CD245 as myosin 18A, a receptor for surfactant A: A novel pathway for activating human NK lymphocytes. *Oncol Immunology* **5**, e1127493 (2016). doi:10.1080/2162402X.2015.1127493 Medline
16. C. H. Yang, J. Szeliga, J. Jordan, S. Faske, Z. Sever-Chroneos, B. Dorsett, R. E. Christian, R. E. Settlege, J. Shabanowitz, D. F. Hunt, J. A. Whitsett, Z. C. Chroneos, Identification of the surfactant protein A receptor 210 as the unconventional myosin 18A. *J. Biol. Chem.* **280**, 34447–34457 (2005). doi:10.1074/jbc.M505229200 Medline
17. F. Petry, M. Botto, R. Holtappels, M. J. Walport, M. Loos, Reconstitution of the complement function in C1q-deficient (C1qa $^{-/-}$) mice with wild-type bone marrow cells. *J. Immunol.* **167**, 4033–4037 (2001). doi:10.4049/jimmunol.167.7.4033 Medline
18. J. Wang, Z.-P. Jiang, N. Su, J.-J. Fan, Y.-P. Ruan, W.-X. Peng, Y.-F. Li, X.-Q. Yu, The role of peritoneal alternatively activated macrophages in the process of peritoneal fibrosis related to peritoneal dialysis. *Int. J. Mol. Sci.* **14**, 10369–10382 (2013). doi:10.3390/ijms140510369 Medline
19. A. Pletinck, R. Vanholder, N. Veys, W. Van Biesen, Protecting the peritoneal membrane: Factors beyond peritoneal dialysis solutions. *Nat. Rev. Nephrol.* **8**, 542–550 (2012). doi:10.1038/nrneph.2012.144 Medline
20. T. Bellón, V. Martínez, B. Lucendo, G. del Peso, M. J. Castro, L. S. Aroeira, A. Rodríguez-Sanz, M. Ossorio, R. Sánchez-Villanueva, R. Selgas, M. A. Bajo, Alternative activation of macrophages in human peritoneum: Implications for peritoneal fibrosis. *Nephrol. Dial. Transplant.* **26**, 2995–3005 (2011). doi:10.1093/ndt/gfq771 Medline
21. J. M. Shipley, R. L. Wesselschmidt, D. K. Kobayashi, T. J. Ley, S. D. Shapiro, Metalloelastase is required for macrophage-mediated proteolysis and matrix invasion in mice. *Proc. Natl. Acad. Sci. U.S.A.* **93**, 3942–3946 (1996). doi:10.1073/pnas.93.9.3942 Medline
22. J. Lu, X. Wu, B. K. Teh, The regulatory roles of C1q. *Immunobiology* **212**, 245–252 (2007). doi:10.1016/j.imbio.2006.11.008 Medline
23. O. R. Colegio, N.-Q. Chu, A. L. Szabo, T. Chu, A. M. Rhebergen, V. Jairam, N. Cyrus, C. E. Brokowski, S. C. Eisenbarth, G. M. Phillips, G. W. Cline, A. J. Phillips, R. Medzhitov, Functional polarization of tumour-associated macrophages by tumour-derived lactic acid. *Nature* **513**, 559–563 (2014). doi:10.1038/nature13490 Medline
24. S. Watanabe, K. Sato, N. Hasegawa, T. Kurihara, K. Matsutani, K. Sanada, T. Hamaoka, S. Fujita, M. Iemitsu, Serum C1q as a novel biomarker of sarcopenia in older adults. *FASEB J.* **29**, 1003–1010 (2015). doi:10.1096/fj.14-262154 Medline
25. C. Shi, P. Velázquez, T. M. Hohl, I. Leiner, M. L. Dustin, E. G. Pamer, Monocyte trafficking to hepatic sites of bacterial infection is chemokine independent and directed by focal intercellular adhesion molecule-1 expression. *J. Immunol.* **184**, 6266–6274 (2010). doi:10.4049/jimmunol.0904160 Medline
26. C. Blériot, T. Dupuis, G. Jouvion, G. Eberl, O. Disson, M. Lecuit, Liver-resident macrophage necroptosis orchestrates type 1 microbicidal inflammation and type-2-mediated tissue repair during bacterial infection. *Immunity* **42**, 145–158

- (2015). [doi:10.1016/j.immuni.2014.12.020](https://doi.org/10.1016/j.immuni.2014.12.020) Medline
27. S. J. Van Dyken, R. M. Locksley, Interleukin-4- and interleukin-13-mediated alternatively activated macrophages: Roles in homeostasis and disease. *Annu. Rev. Immunol.* **31**, 317–343 (2013). [doi:10.1146/annurev-immunol-032712-095906](https://doi.org/10.1146/annurev-immunol-032712-095906) Medline
 28. A. J. Byrne, T. M. Maher, C. M. Lloyd, Pulmonary Macrophages: A New Therapeutic Pathway in Fibrosing Lung Disease? *Trends Mol. Med.* **22**, 303–316 (2016). [doi:10.1016/j.molmed.2016.02.004](https://doi.org/10.1016/j.molmed.2016.02.004) Medline
 29. J. Xue, V. Sharma, M. H. Hsieh, A. Chawla, R. Murali, S. J. Pandol, A. Habtezion, Alternatively activated macrophages promote pancreatic fibrosis in chronic pancreatitis. *Nat. Commun.* **6**, 7158 (2015). [doi:10.1038/ncomms8158](https://doi.org/10.1038/ncomms8158) Medline
 30. J. M. Coya, H. T. Akinbi, A. Sáenz, L. Yang, T. E. Weaver, C. Casals, Natural anti-infective pulmonary proteins: In vivo cooperative action of surfactant protein SP-A and the lung antimicrobial peptide SP-BN. *J. Immunol.* **195**, 1628–1636 (2015). [doi:10.4049/jimmunol.1500778](https://doi.org/10.4049/jimmunol.1500778) Medline
 31. I. García-Verdugo, F. Sánchez-Barbero, F. U. Bosch, W. Steinhilber, C. Casals, Effect of hydroxylation and N187-linked glycosylation on molecular and functional properties of recombinant human surfactant protein A. *Biochemistry* **42**, 9532–9542 (2003). [doi:10.1021/bi0347196](https://doi.org/10.1021/bi0347196) Medline
 32. F. Sánchez-Barbero, G. Rivas, W. Steinhilber, C. Casals, Structural and functional differences among human surfactant proteins SP-A1, SP-A2 and co-expressed SP-A1/SP-A2: Role of supratrimeric oligomerization. *Biochem. J.* **406**, 479–489 (2007). [doi:10.1042/BJ20070275](https://doi.org/10.1042/BJ20070275) Medline
 33. F. Sánchez-Barbero, J. Strassner, R. García-Cañero, W. Steinhilber, C. Casals, Role of the degree of oligomerization in the structure and function of human surfactant protein A. *J. Biol. Chem.* **280**, 7659–7670 (2005). [doi:10.1074/jbc.M410266200](https://doi.org/10.1074/jbc.M410266200) Medline
 34. G. Li, J. Siddiqui, M. Hendry, J. Akiyama, J. Edmondson, C. Brown, L. Allen, S. Levitt, F. Poulain, S. Hawgood, Surfactant protein-A-deficient mice display an exaggerated early inflammatory response to a beta-resistant strain of influenza A virus. *Am. J. Respir. Cell Mol. Biol.* **26**, 277–282 (2002). [doi:10.1165/ajrcmb.26.3.4584](https://doi.org/10.1165/ajrcmb.26.3.4584) Medline
 35. F. Carlucci, A. Ishaque, G. S. Ling, M. Szajna, A. Sandison, P. Donatien, H. T. Cook, M. Botto, C1q Modulates the Response to TLR7 Stimulation by Pristane-Primed Macrophages: Implications for Pristane-Induced Lupus. *J. Immunol.* **196**, 1488–1494 (2016). [doi:10.4049/jimmunol.1401009](https://doi.org/10.4049/jimmunol.1401009) Medline
 36. D. R. Herbert, C. Hölscher, M. Mohrs, B. Arendse, A. Schwegmann, M. Radwanska, M. Leeto, R. Kirsch, P. Hall, H. Mossmann, B. Claussen, I. Förster, F. Brombacher, Alternative macrophage activation is essential for survival during schistosomiasis and downmodulates T helper 1 responses and immunopathology. *Immunity* **20**, 623–635 (2004). [doi:10.1016/S1074-7613\(04\)00107-4](https://doi.org/10.1016/S1074-7613(04)00107-4) Medline
 37. A. A. Thompson, P. M. Elks, H. M. Marriott, S. Eamsamrנג, K. R. Higgins, A. Lewis, L. Williams, S. Parmar, G. Shaw, E. E. McGrath, F. Formenti, F. J. Van Eeden, V. L. Kinnula, C. W. Pugh, I. Sabroe, D. H. Dockrell, E. R. Chilvers, P. A. Robbins, M. J. Percy, M. C. Simon, R. S. Johnson, S. A. Renshaw, M. K. B. Whyte, S. R. Walmsley, Hypoxia-inducible factor 2 α regulates key neutrophil functions in humans, mice, and zebrafish. *Blood* **123**, 366–376 (2014). [doi:10.1182/blood-2013-05-500207](https://doi.org/10.1182/blood-2013-05-500207) Medline
 38. S. J. Jenkins, D. Ruckerl, P. C. Cook, L. H. Jones, F. D. Finkelman, N. van Rooijen, A. S. MacDonald, J. E. Allen, Local macrophage proliferation, rather than recruitment from the blood, is a signature of TH2 inflammation. *Science* **332**, 1284–1288 (2011). [doi:10.1126/science.1204351](https://doi.org/10.1126/science.1204351) Medline
 39. S. J. Jenkins, D. Ruckerl, G. D. Thomas, J. P. Hewitson, S. Duncan, F. Brombacher, R. M. Maizels, D. A. Hume, J. E. Allen, IL-4 directly signals tissue-resident macrophages to proliferate beyond homeostatic levels controlled by CSF-1. *J. Exp. Med.* **210**, 2477–2491 (2013). [doi:10.1084/jem.20121999](https://doi.org/10.1084/jem.20121999) Medline
 40. L. H. Jackson-Jones, D. Ruckerl, F. Svedberg, S. Duncan, R. M. Maizels, T. E. Sutherland, S. J. Jenkins, H. J. McSorley, C. Bénézec, A. S. MacDonald, J. E. Allen, IL-33 delivery induces serous cavity macrophage proliferation independent of interleukin-4 receptor alpha. *Eur. J. Immunol.* **46**, 2311–2321 (2016). [doi:10.1002/eji.201646442](https://doi.org/10.1002/eji.201646442) Medline
 41. M. Camberis, G. Le Gros, J. Urban Jr., Animal model of *Nippostrongylus brasiliensis* and *Heligmosomoides polygyrus*. *Curr. Protoc. Immunol.* **Chapter 19**, 12 (2003). [doi:10.1002/0471142735.im1912s55](https://doi.org/10.1002/0471142735.im1912s55) Medline
 42. M. J. Holland, Y. M. Harcus, P. L. Riches, R. M. Maizels, Proteins secreted by the parasitic nematode *Nippostrongylus brasiliensis* act as adjuvants for Th2 responses. *Eur. J. Immunol.* **30**, 1977–1987 (2000). [doi:10.1002/1521-4141\(200007\)30:7<1977::AID-IMMU1977>3.0.CO;2-3](https://doi.org/10.1002/1521-4141(200007)30:7<1977::AID-IMMU1977>3.0.CO;2-3) Medline
 43. T. E. Sutherland, N. Logan, D. Ruckerl, A. A. Humbles, S. M. Allan, V. Papayannopoulos, B. Stockinger, R. M. Maizels, J. E. Allen, Chitinase-like proteins promote IL-17-mediated neutrophilia in a tradeoff between nematode killing and host damage. *Nat. Immunol.* **15**, 1116–1125 (2014). [doi:10.1038/ni.3023](https://doi.org/10.1038/ni.3023) Medline
 44. C. Casals, J. Arias-Díaz, F. Valiño, A. Sáenz, C. García, J. L. Balibrea, E. Vara, Surfactant strengthens the inhibitory effect of C-reactive protein on human lung macrophage cytokine release. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **284**, L466–L472 (2003). [doi:10.1152/ajplung.00325.2002](https://doi.org/10.1152/ajplung.00325.2002) Medline
 45. W. Q. Zeng, J. Q. Zhang, Y. Li, K. Yang, Y. P. Chen, Z.-J. Liu, A new method to isolate and culture rat kupffer cells. *PLOS ONE* **8**, e70832 (2013). [doi:10.1371/journal.pone.0070832](https://doi.org/10.1371/journal.pone.0070832) Medline
 46. A. C. MacKinnon, S. L. Farnworth, P. S. Hodgkinson, N. C. Henderson, K. M. Atkinson, H. Leffler, U. J. Nilsson, C. Haslett, S. J. Forbes, T. Sethi, Regulation of alternative macrophage activation by galectin-3. *J. Immunol.* **180**, 2650–2658 (2008). [doi:10.4049/jimmunol.180.4.2650](https://doi.org/10.4049/jimmunol.180.4.2650) Medline

ACKNOWLEDGMENTS

The authors thank M. Waterfall for expertise with flow cytometry; M. Botto and M. Gray for providing C1q^{-/-} mice; C. Benezech for providing IL-4R α ^{-/-} mice; S. Walmsley for providing LysM^{Cre/WT} / HIF1 α ^{fl/fl} mice; D. Ruckerl, M. Lorente, R. Tillu, for valuable advice; G. Goodman and M. Rigol-Muxart for veterinary advice; S. Jenkins for critical evaluation of the manuscript; A. Fulton and S. Duncan for excellent technical assistance, and support staff for excellent animal husbandry. The Myo18 neutralising antibody is covered by an MTA between Edinburgh University and Pennsylvania State University (PSU). The antibody was provided by Dr Zissis C. Chronos from PSU, who is an author on this paper. C.M. Minutti was recipient of fellowships from the Spanish Ministry of Science (FPU-AP2010-1524 and Est13/00372) and Institute of Health Carlos III (CIBERES). E. Ringqvist gratefully acknowledges funding support from the Wenner-Gren Foundation. This work was supported by the Spanish Ministry of Economy and Competitiveness (SAF2012-32728 and SAF2015-65307-R) and Institute of Health Carlos III (CIBERES-CB06/06/0002) to C. C., Medical Research Council UK (MR/K01207X/1) and Wellcome Centre for Cell-Matrix Research support to J. E. A., Medical Research Council (MR/M011755/1) and European Union (CIG-631413) to D.M.Z., National of Institute Health grants HL068127 and HL128746 to Z. C. German Research Council (DFG) grant 609/2-1 to C.S., a Wellcome Trust Intermediate Fellowship to D.F. and a Wenner Gren Foundation Fellowship to E.R. All data to understand and assess the conclusions of this research are available in the main text or supplementary materials. C.M.M. designed and performed research, analyzed and interpreted data and wrote the manuscript; L.H.J.-J., B.G.F., T.E.S., J.A.K., N.L. and E.R. performed research, contributed to experimental design and manuscript preparation; R.G-P. performed research; D.A.F., A.A., C.S. and Z.C. contributed tools, provided expertise and edited the manuscript; D.M.Z. contributed to project funding, provided expertise and edited the manuscript; C.C. and J.E.A. funded and designed the research, organized the project and analyses, and wrote the manuscript. All authors reviewed and approved the final version of the manuscript. The authors declare no conflict of interest.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/cgi/content/full/science.aaj2067/DC1

Materials and Methods

Figs. S1 to S14

Tables S1 and S2

References (30–46)

13 September 2016; resubmitted 11 March 2017

Published online 11 May 2017

10.1126/science.aaj2067




Fig. 1. Higher worm burden, greater nematode-induced lung damage and reduced IL-4-induced proliferation and activation in mice lacking SP-A. Samples were assessed 6 days after *N. brasiliensis* (Nb) infection. (A) SP-A protein expression in lung tissue of WT and IL-4R α ^{-/-} mice. (B) Adult larvae in the small intestine. (C) Lung damage, quantified by 'mean linear intercept' from micrographs of H&E stained lung sections, and (D) microscopy of H&E stained lung sections (scale bars, 500 μ m) in WT and SP-A^{-/-} mice. (E) Amplification of *Col1a1*- and (F) *Mmp12*-encoding mRNA in lung tissue. Number of (G) red blood cells and (H) neutrophils isolated in BAL. Expression of (I) RELM α by aM ϕ from BAL. (J) BrdU incorporation and (K) Ki67 expression by aM ϕ from BAL. Data are representative from two independent experiments (mean \pm SEM; naive: 3 mice, Nb: 6 mice). (L to N) WT and SP-A^{-/-} mice treated with 5 μ g IL-4c (i.p.) at days 0 and 2 and analyzed at day 4. (L) Relative SP-A levels in BAL (representative western blot shown) of WT mice treated with IL-4c or PBS. (M) RELM α expression and (N) BrdU incorporation in aM ϕ . Data pooled from three independent experiments (means \pm SEM) (PBS: 9 mice, IL-4c: 11 mice). ANOVA followed by the Bonferroni multiple-comparison test was used. * p < 0.05, ** p < 0.01, and *** p < 0.001 when compared with the untreated/uninfected group. $^{\circ}p$ < 0.05, $^{\circ\circ}p$ < 0.01 and $^{\circ\circ\circ}p$ < 0.001 when WT vs. SP-A^{-/-} groups are compared.




Fig. 2. SP-A and C1q act through Myo18A to enhance IL-4-induced proliferation and activation of alveolar and peritoneal macrophages, respectively. (A) Murine macrophages were treated with IL-4 in the presence or absence of SP-A or C1q. BrdU incorporation and Ym1 secretion are shown. (B) For aMφs (red), 5 μ g IL-4c was delivered ip at days 0 and 2, and BAL cells analyzed at day 4. For pMφs (blue), 1 μ g IL-4c was delivered ip at day 0, and peritoneal cells analyzed at day 1; BrdU incorporation and RELM α expression are shown. (C) Murine macrophages were treated with anti-Myo18A or rabbit IgG plus either IL-4+SP-A (aMφ) or IL-4+C1q (pMφ). BrdU incorporation and Ym1 secretion are shown. (D) Concurrently with IL-4c delivery, some WT mice were treated intra-nasally or ip with either anti-Myo18A or rabbit IgG antibody. BrdU incorporation and RELM α expression are shown in aMφs and pMφs. All statistical analysis was performed by ANOVA followed by the Bonferroni multiple-comparison test. (A and C) Results are presented as means (\pm SEM) from three different cell cultures with at least three biological replicates. * p < 0.05, ** p < 0.01, and *** p < 0.001, when compared with untreated cells; $^{\circ}p$ < 0.05, $^{\circ\circ}p$ < 0.01, and $^{\circ\circ\circ}p$ < 0.001, when SP-A+IL-4- or C1q+IL-4- treated are compared with IL-4-treated; ### p < 0.01, and ### p < 0.001, the effect of anti-Myo18A antibody on cells treated with SP-A+IL-4 or C1q+IL-4. (B and D) Data were pooled from three independent experiments (means \pm SEM) (PBS: 6 mice, other groups: 9 mice). * p < 0.05, ** p < 0.01, and *** p < 0.001, when compared with PBS treated mice; $^{\circ}p$ < 0.05, when WT vs. C1q $^{-/-}$ mice treated with IL-4c are compared (B); $^{\circ}p$ < 0.05, $^{\circ\circ}p$ < 0.01, and $^{\circ\circ\circ}p$ < 0.001 when anti-Myo18A vs rabbit IgG treatment is compared in IL-4c-treated mice (D).




Fig. 3. C1q enhances peritoneal fibrosis induced by a lactate dialysate. WT, C1qa^{-/-} or IL-4Rα^{-/-} mice were either untreated (C) or injected ip with Dianeal PD-4 every other day for 28 days. Samples were analyzed a day after the last delivery. (A) Total amount of C1q in the peritoneal washes was determined by ELISA. (B) Quantification of the thickness of the submesothelial compact zone from (C) microscopy of Masson's trichrome stained parietal peritoneum slices (scale bars, 0.1 mm). Amplification of (D) *Col1a1*- and (E) *Acta2*- encoding mRNA in peritoneal tissue. Expression of (F) RELMα, (G) Ym1, (H) Arg and (I) Ki67 by pMφs. Results are representative from two independent experiments (means ± SEM) (untreated: 3 mice, PD4: 6 mice). ANOVA followed by the Bonferroni multiple-comparison test or Student's *t*-test (A) was used. **p* < 0.05, ***p* < 0.01, and ****p* < 0.001 when compared with control group; °*p* < 0.05, °°*p* < 0.01, and °°°*p* < 0.001 when WT vs. C1qa^{-/-} mice treated with Dianeal PD-4 are compared..




Fig. 4. C1q is required for appropriate macrophage activation in the liver during *Listeria monocytogenes* infection. (A to C) WT or C1qa^{-/-} mice received 1 μg IL-4c (ip) at day 0, and samples were analyzed at day 1. (A) Myo18A expression on the surface of resident macrophages (identified as described in the methods) from the indicated tissues. (B) BrdU incorporation and RELMα expression of liver macrophages. (C) (upper panel) IL-4-induced amplification of C1q-encoding mRNA in the liver. (C to I) WT, C1qa^{-/-}, or IL-4Rα^{-/-} mice were left uninfected or received intravenous infection with 10⁴ *L. monocytogenes* (Lm) c.f.u., and samples were assessed at 3.5 days p.i. (C) (lower panel) *L. monocytogenes*-induced amplification of C1q-encoding mRNA. (D) Expression of RELMα by liver macrophages. (E) BrdU incorporation by liver macrophages. (F) Quantification of ALT and AST in serum. (G) Amplification of Acta2- and Col1a1-encoding mRNA in the liver. (H) Liver bacterial load. (I) Number of monocytes in liver single cell suspensions. Data are representative from two independent experiments (mean ± SEM; naïve: 4 mice, Lm: 5 mice). ANOVA followed by the Bonferroni multiple-comparison test was used. **p* < 0.05, and ****p* < 0.001, when compared with the uninfected group; °*p* < 0.05, °°*p* < 0.01, and °°°*p* < 0.001 when WT vs. C1qa^{-/-} or IL-4Rα^{-/-} infected groups are compared.



Local amplifiers of IL-4R α -mediated macrophage activation promote repair in lung and liver

Carlos M. Minutti, Lucy H. Jackson-Jones, Belén García-Fojeda, Johanna A. Knipper, Tara E. Sutherland, Nicola Logan, Emma Rinqvist, Raquel Guillamat-Prats, David A. Ferenbach, Antonio Artigas, Cordula Stamme, Zissis C. Chroneos, Dietmar M. Zaiss, Cristina Casals and Judith E. Allen (May 11, 2017)
published online May 11, 2017

Editor's Summary

This copy is for your personal, non-commercial use only.

Article Tools Visit the online version of this article to access the personalization and article tools:
<http://science.sciencemag.org/content/early/2017/05/10/science.aaj2067>

Permissions Obtain information about reproducing this article:
<http://www.sciencemag.org/about/permissions.dtl>

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published weekly, except the last week in December, by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. Copyright 2016 by the American Association for the Advancement of Science; all rights reserved. The title *Science* is a registered trademark of AAAS.